
200600228

No.



HHIR UNIVERD STRAFFS OF ANTERIO

<u> Thiese; Presents; Shalil, Come;</u>

Enzn Znden Beheer B. A.

HICCOS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OF ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY FEARS FROM THE DATE OF THIS GRAND, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC **REPLENISHMENT** OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY ${
m LAW}$, THE GENERO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR MING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE URPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT (BY THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

LETTUCE

'Tellmark'

In Testimony Whereat, I have hereunto set my hand and caused the seal of the Plant Buriety Frotestion Office to be affixed at the City of Washington, D.C. this fifth day of June, in the year two thousand and eight.

Plant Variety Protection Office Agricultural Marketing Service

Colward T. Schafer

(See reverse for instructions and information collection burden statement)

DATE

CAPACITY OR TITLE

06/05/2006

V. LAMBALE

HIRECTOR

GENERAL INSTRUCTIONS: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner; (2) completed exhibits A, B, C, E, F; (3) for a tuber reproduced variety, verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; and (4) payment by credit card or check drawn on a U.S. bank for \$4,382 (\$518 filing fee and \$3,864 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice). NEW: With the application for a seed reproduced variety or by direct deposit soon after filing, the applicant must provide at least 3,000 viable untreated seeds of the variety per se, and for a hybrid variety at least 3,000 untreated seeds of each line necessary to reproduce the variety. Partial applications will be held in the PVPO for not more than 90 days; then returned to the applicant as un-filed. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a payment by credit card or check payable to "Treasurer of the United States" in the amount of \$768 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. The fees for filing a change of address; owner's representative; ownership or assignment; or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

Plant Variety Protection Office

Telephone: (301) 504-5518

FAX: (301) 504-5291

General E-mail: PVPOmail@usda.gov

Homepage: http://www.ams.usda.gov/science/pvpo/PVPindex.htm

SPECIFIC INSTRUCTIONS:

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority and **provide evidence** that the permanent name of the application variety (even if it is a parental, inbred line) has been cleared by the appropriate recognized authority before the Certificate of Protection is issued. For example, for agricultural and vegetable crops, contact: U.S. Department of Agriculture, Agricultural Marketing Service, Livestock and Seed Programs, **Seed Regulatory and Testing Branch**, 801 Summit Crossing Place, Suite C, Gastonia, North Carolina 28054-2193 Telephone: (704) 810-8870. http://www.ams.usda.gov/lsg/seed.htm.

ITEM

19a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;

(3) evidence of uniformity and stability; and

- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 19b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach replicated statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 19c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 19d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 19e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 20. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 23. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 24. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 22. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)
- 23. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

FIRST DATE OF SALES IN US: 06/10/2005

24. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

variety is protected by intellectual property right (Plant Breeder's Right or Patent).)
HALLMARK, PVP US, #9800266, DATE ISSUED: 09/25/2003

TELLURIDE, P.V.P. US, #200300168, DATE 1850ED: 07/26/2005

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

Exhibit A: Origin and Breeding History of 'Tellmark', (CS14.2063)

'<u>Tellmark'</u>, (CS14.2063)

Female Male 'Hallmark'-----X----- 'Telluride'- (August 1999)

F₁--- Twenty four (24) seeds of the cross were tested with *Bremia lactucae*, pathotype CA-V. An eight (8) plant mass and three (3) single plant selections were made for a seed increase. (February 2000)

F₂--- Seeds from the mass and single plant selection were sown (August 2000) on corky root (*Sphingomonas suberifaciens*) infested test plots in Salinas, California. Eight (8) plants from the mass were selected (October 2000) and transferred to the greenhouse at the company facility in San Juan Bautista, California for seed increase.

F₃---Selected plants were tested for resistance to *Bremia lactucae*, pathotype CA-V and Twelve (12) single plant selections were saved for seed increase.(March 2001)

F₄--- Seeds of the selections above were sown (August 2001) on a corky root infested test plot in Salinas, California. Sixteen (16) single plant selections were made (October 2001) and transferred to the company greenhouses in San Juan Bautista, California for seed increase.

F₅--- Seed from the selected plants were sown at the company trial grounds in San Juan Bautista, California (May 2002). One line from the seed increase above looked uniform for type with better size and type and 8 single plant selections were made for seed increase at the company facilities.

F₆--- Seed from six (6) of the single plant selections were sown at leased trial plots at Griffith, NSW, Australia (October 2002). Two lines were uniform for size, type and maturity. Twenty-five (25) plants of each line were selected for seed increase and trials. The two lines were tested in 2003 as CS14.2063 and CS14.2064. Seed of both lines were tested and found to be resistant for *Bremia lactucae*, pathotype CA-V. Molecular marker tests were conducted and confirmed the presence of the marker for the resistance factor to *Bremia lactucae*, pathotypes, CA-I through CA-VI. Bioassays and molecular marker tests confirmed resistance to corky root (*Sphingomonas suberifaciens*).

F₇--- Trials of CS14.2063 were conducted at the company trial facilities and commercial lettuce production fields in the Salinas Valley of California. CS14.2063 was selected for seed increase for larger commercial trials. Twenty five plants (25) were selected at the company facility in San Juan Bautista, California for foundation seed (June 2003). The foundation seed was tested with bioassays and molecular markers to confirm resistance to corky root and *Bremia lactucae*, pathotypes CA-I through CA-VI.

(revised)

 F_8 ---The foundation seed was sown in October 2003 for advanced commercial trials and sales at Hillston, NSW, Australia and rogued for type by the research staff.

The breeding method employed was pedigree selection, using both single plant selection and mass selection practices. The selection criteria for cv. 'Tellmark' were:

- 1. A cultivar with uniformity larger frame, head diameter and shorter core when compared to 'Telluride'
- 2. Resistance to *Bremia lactucae*, pathotype, CA-I through CA-VI where 'Hallmark' is susceptible to CA-V and CA-VI.

In three cycles of seed increase we have noted a large non-heading, dark green variant at the rate of one plant per 5,000 plants. This is well within the range of variants noted for most 'Salinas- type' icebergs. After two generations of reproduction we believe this variety is uniform and stable for type.

Exhibit B: Novelty Statement of Lettuce cv. 'Tellmark', (CS14.2063)

'Tellmark' is a vigorous, crisphead lettuce cultivar adapted to Coastal California. Its optimum sowing period is from February through August in Coastal California. 'Tellmark' has black seed and is resistant to *Bremia lactucae*, pathotypes CA-I through CA-VI (Tables 2a, b, 3). 'Tellmark' is resistant to corky root (*Sphingomonas suberifaciens*) and contains the gene *cor* (Tables 4, 5).

'Tellmark' is phenotypically distinct from its most similar commercial cultivar, 'Telluride'. 'Tellmark' has a significantly larger frame diameter at the 95% confidence level with means ranging from 52.2 to 62.6cm compared to 'Telluride' with means ranging from 48.8 to 54.5cm in 4 of 4 tests(Table 1). 'Tellmark' has a significantly larger head diameter at the 95% confidence level with means ranging from 12.8 to 15.9 cm compared to 'Telluride' with means ranging from 12.8 to 14.1cm in 3 of 4 tests (Table 1). 'Tellmark' has a significantly shorter core length at the 95% confidence level with means ranging from 23.8 to 38.4mm compared to 'Telluride' with means ranging from 28.3 to 46.9mm in 3 of 4 tests (Table 1). 'Tellmark' is lighter green in color than 'Telluride' 137C vs. 138A respectively (RHS color chart). The head shape of 'Tellmark' is slightly flattened while the head shape of 'Telluride' is spherical. Leaf blistering is absent/slight on 'Tellmark' while moderate on 'Telluride'.

'Tellmark' is resistant to *Bremia lactucae*, pathotypes CA-V and CA-VI whereas the cultivar 'Hallmark' is susceptible (Tables 2a, b, 3).

Exhibit C

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 1.4 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY **PLANT VARIETY PROTECTION OFFICE** BELTSVILLE, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY Lettuce (Lactuca sativa L.)

	-	
NAME OF APPLICANT (S) ENZA ZADEN BEHEER B.V.	TEMPORARY OR EXPERIMENTAL DESIGNATION 14. 2063	VARIETY NAME TELLMARK
ADDRESS (Street and No. or RD No., City, State, Zip Code, and Country) POST BUS 7, 1600 AA ENKHUIZEN HALING 1 ^E , 1602 DB ENKHUIZEN THE NETHERLANDS		PVPO NUMBER 200600228
Place the appropriate number that describes the varietal is either 99 or less or 9 or less. Measured data should be recognized color standard may be used to determine pla	e the mean of an appropriate number (at least 20) o	
The Location of the Test Area is: SAN JUAN BAUTISTA, SALIMAS, SANTA P	Color System Used:	RHS
SPECIFIC VARIETIES USED FOR COMPARISON AS Compounding to the comparison varieties must be the compounding to	CHECK VARIETIES IN THIS APPLICATION: Use s most similar variety used in Exhibit B. Most Similar Variety (c1)	tandard regional check varieties, which are adapted to
1. PLANT TYPE: (See List of Suggested Check Varieties 01 = Cutting/Leaf	naine 07 = Salinas Group 10 = L s Group 08 = Eastern (Ithaca) Group 11 = C	atin Other (Specify) 0 7
2. SEED: (a1) 2 (c1) 2 COLOR (a1) 1 = White (Silver Gray) 2 = Black (Grey Brown) (c1) 3 = Brown (Amber) (c2)	LIGHT DORMANCY (a1) 1 = Light Required 2 = Light Not Required (c1) (c2)	The state of the
3. COTYLEDON TO FOURTH LEAF STAGE: NOTE: P condition		h leaf from 20 day-old seedling grown under optimal
SHAPE OF COTYLEDONS: 1 = Broad (a1)	$ \begin{array}{ccc} 2 & = & & & & & & & & & \\ 2 & & & & & & & \\ \hline 2 & & & & & & \\ \end{array} $ (c1)	(c2) 2
SHAPE OF FOURTH LEAF: (a1)	(c1) 4	(c2) 3

3. COTYLEDON TO FOURTH LEAF STAGE: (continued)







2. Round



3. Oval

4. Elongated



5. Lanceolate



6. Pinnately lobed

LENGTH/WIDTH INDEX OF FOURTH LEAF: L/W x 10

1 3 (a1)

14

APICAL MARGIN:

1 = Entire

2 = Crenate/Gnawed

4 = Moderately Dentate 5 = Coarsely Dentate

7 = Lobed

8 = Other (Specify)

3 = Finely Dentate 6 = Incised

(a1)

(c1)

BASAL MARGIN: (Use the options for Apical Margin above)

(a1)

(c1)

UNDULATION:

1 = Flat

2 = Slight

3 = Medium

4 = Marked

(c1)

2

3

GREEN COLOR: 1 = Yellow Green

2 = Light Green

3 = Medium Green

5 = Blue Green

7 = Grey Green

4 = Dark Green

6 = Silver Green

(a1)

(a1)

3

2

(c1)

3

ANTHOCYANIN:

DISTRIBUTION:

1 = Absent

2 = Margin Only

3 = Spotted

4 = Throughout

5 = Other (Specify)

(a1)

į

(c1)

CONCENTRATION:

1 = Light

2 = Moderate

3 = Intense

(a1)

(a1)

(a1)

(c1)

ROLLING:

1 = Absent

2 = Present

(c1)

CUPPING:

1 = Uncupped

2 = Slight

(c1)

1

3 = Markedly

(c2)

REFLEXING:

1 = None

2 = Apical Margin

3 = Lateral Margins

(a1)

(c1)

(c2)

4. MATURE LEAVES (Observe Harvest-Mature Outer Leaves)

NOTE: Provide color photo of a harvest-mature leaf which accurately shows color and margin characteristics.

MARGIN:

INCISION DEPTH:

1 = Absent/Shallow (Dark Green Boston)

2 = Moderate (Vanguard)

3 = Deep (Great Lakes 659)

(deepest penetration of the margin)

(a1)

(c1)

INDENTATION: (Finest divisions of the margin)

1 = Entire (Dark Green Boston)

2 = Shallowly Dentate (Great Lakes 65)

4 = Crenate (Vanguard) 5 = Other (Specify)

3 = Deeply Dentate (Great Lakes 659)

(a1)

4 (c1)

2 (c2)

UNDULATIONS OF THE APICAL MARGIN:

1 = Absent/Slight (Dark Green Boston) 2 = Moderate (Vanguard)

3 = Strong (Great Lakes 659)

2 (a1)

2 (c1)

(c2)

GREEN COLOR:

1 = Very Light Green (Bibb)

(a1)

3 = Medium Green (Great Lakes)

5 = Very Dark Green 6 = Other (Specify)

2 = Light Green (Minetto) 4 = Dark Green (Vanguard)

(c1)

3 (c2)

ANTHOCYANIN:

DISTRIBUTION:

1 = Absent

2 = Margin Only (Big Boston)

3 = Spotted (California Cream Butter)

5 = Other (Specify)

4 = Throughout (Prize Head)

(a1)

(c1)

(c2)

CONCENTRATION:

1 = Light (Iceberg)

2 = Moderate (Prize Head) 3 = Intense (Ruby)

(a1)

(c1)

(c1)

2

(c2)

SIZE:

1 = Small

2= Medium

3 = Large

3

GLOSSINESS:

1 = Dull (Vanguard)

(a1)

(a1)

(a1)

(a1)

2 = Moderate (Salinas)

3 = Glossy (Great Lakes)

(c2)

(a1)

2

0

BLISTERING:

1 = Absent/Slight

2 = Moderate

3

(c1)

(Salinas)

0 3

(Vanguard)

(Prize Head)

3 = Strong

0

LEAF THICKNESS:

1 = Thin

2 = Intermediate

2 (c1) 0

3 = Thick

D 2

TRICHOMES:

1 = Absent (Smooth)

2 = Present (Spiny)

(c1)

5. PLANT:

SPREAD OF FRAME LEAVES:

(a1)

HEAD BIAMPTER: (Masket Trimmed with Single Cap Lear)	5. PLANT: (continued)							
HEAD SHAPE: 1 = Flattened 2 = Slightly Flattened 4 + Elongale 5 = Nor-Heading 5 = Nor-H	HEAD DIAMETER: (Market Trime	med with Single Cap I	Leaf)					
6 = Other (Specify)		(a1)	14 cm	(c1)	/ 3 cm	(c2)	1 4 cm	
Call O D D Call O D Call O D D Call O D D Call O D D Call O D D D Call O D D D Call O D D D Call O D D D D D D D D D	HEAD SHAPE:		3 = Sphe	erical	5 = Non-Heading			
HEAD SIZE CLASS: 1 = Small	6 = Other (Specify)	2 = Slightly Fla	ittened 4 = Elon 	gate				
(a1) 0 3 (c1) 0 2 (c2) 0 3		(a1)	02	(c1)	03	(c2)	03	
HEAD PER CARTON: (a1)	HEAD SIZE CLASS:	1 = Small	2 = Medi	ium	3= Large			
HEAD WEIGHT:		(a1)	0 3	(c1)	02	(c2)	03	
HEAD WEIGHT:	HEAD PER CARTON:							
HEAD FIRMNESS: 1 = Loose		(a1)	24	(c1)	24	(c2)	24	
HEAD FIRMNESS: 1 = Loose 2 = Moderate 3 = Firm 4 = Very Firm				(0.7	<u> </u>	(02)		
HEAD FIRMNESS: 1 = Loose 2 = Moderate 3 = Firm 4 = Very Firm	HEAD WEIGHT:		9					
6. BUTT: SHAPE: 1 = Slightly Concave (a1) 2		(a1)	1/14	g. (c1)	6 6 4	g. (c2)	691	g.
6. BUTT: SHAPE: 1 = Slightly Concave (a1) 2	HEAD FIRMNESS:	1 = Loose	2 = Moderate		3= Firm	4	= Very Firm	
SHAPE: 1 = Slightly Concave (a1) 2 (c1) 2 (c2) 2 MIDRIB: 1 = Flattened (Salinas) (a1) 1 (c1) 2 (c2) 2 MIDRIB: 1 = Flattened (Salinas) (a1) 1 (c1) 1 (c2) 1 7. CORE: DIAMETER AT BASE OF HEAD: (a1) 3 2 mm (c1) 3 2 mm (c2) 3 1 mm RATIO OF HEAD DIAMETER: (a1) 4 5 (c1) 4 0 (c2) 4 3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: O5/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 3 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:		(a1)	2	(c1)	3	(c2)	3	
SHAPE: 1 = Slightly Concave (a1) 2 (c1) 2 (c2) 2 MIDRIB: 1 = Flattened (Salinas) (a1) 1 (c1) 2 (c2) 2 MIDRIB: 1 = Flattened (Salinas) (a1) 1 (c1) 1 (c2) 1 7. CORE: DIAMETER AT BASE OF HEAD: (a1) 3 2 mm (c1) 3 2 mm (c2) 3 1 mm RATIO OF HEAD DIAMETER: (a1) 4 5 (c1) 4 0 (c2) 4 3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: O5/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 3 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	6. BUTT:		, , , , , , , , , , , , , , , , , , ,					
MIDRIB: 1 = Flattened (Salinas) 2 = Moderately Raised 3 = Prominently Raised (Great Lakes 659)	•	1 = Slightly Cor	ncave 2 =	Flat :	3 = Rounded			
MIDRIE: 1 = Flattened (Salinas) 2 = Moderately Raised 3 = Prominently Raised (Great Lakes 659) (c1) 1 (c2) 1 7. CORE: DIAMETER AT BASE OF HEAD: (a1) 3 2 mm (c1) 3 2 mm (c2) 3 1 mm RATIO OF HEAD DIAMETER: (a1) 4 5 (c1) 4 0 (c2) 4 3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 2 = Slow 3 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:						(c2)	2	
(a1) 1 (c1) 1 (c2) 1 7. CORE: DIAMETER AT BASE OF HEAD: (a1) 3 2 mm (c1) 3 2 mm (c2) 3 1 mm RATIO OF HEAD DIAMETER: (a1) 4 5 (c1) 4 0 (c2) 4 3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 3 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:					<u> </u>			
7. CORE: DIAMETER AT BASE OF HEAD: (a1) 32 mm (c1) 32 mm (c2) 31 mm RATIO OF HEAD DIAMETER: (a1) 4.5 (c1) 4.0 (c2) 4.3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 39 mm (c1) 42 mm (c2) 37 mm 8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 96 (c1) 93 (c2) 91 BOLTING CLASS: 1 = Very Slow 2 = Medium 2 = Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	MIDRIB:				[ed (Great Lakes 6	59)
DIAMETER AT BASE OF HEAD: (a1) 3 2 mm (c1) 3 2 mm (c2) 3 1 mm RATIO OF HEAD DIAMETER/CORE DIAMETER: (a1) 4 5 (c1) 4 0 (c2) 4 3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 3 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:		(a1)		(c1)		(c2)		
RATIO OF HEAD DIAMETER/CORE DIAMETER: (a1) 4,5 (c1) 4,0 (c2) 4,3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: 05/24/2 005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	7. CORE:							
RATIO OF HEAD DIAMETER/CORE DIAMETER: (a1) 4 5 (c1) 4 0 (c2) 4 3 CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 3 9 mm (c1) 4 2 mm (c2) 3 7 mm 8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 9 6 (c1) 9 3 (c2) 9 1 BOLTING CLASS: 1 = Very Slow 3 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	DIAMETER AT BASE OF HE		20	(-4)		(
CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 39 mm (c1) 42 mm (c2) 37 mm 8. BOLTING: (Give First Water Date: 05/24/2 005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 96 (c1) 93 (c2) 91 BOLTING CLASS: 1 = Very Slow 2 = Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:		(a1)	[3 2 mm	(C1)	3 2 mm	(c2)	3 1 mm	
CORE HEIGHT FROM BASE OF HEAD TO APEX: (a1) 39 mm (c1) 42 mm (c2) 37 mm 8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 96 (c1) 93 (c2) 91 BOLTING CLASS: 1 = Very Slow 2 = Medium 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	RATIO OF HEAD DIAMETER	CORE DIAMETER:						
(a1) 39 mm (c1) 42 mm (c2) 37 mm 8. BOLTING: (Give First Water Date: 05/24/2 005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 96 (c1) 93 (c2) 91 BOLTING CLASS: 1 = Very Slow 3 = Medium 5 = Very Rapid 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	•	(a1)	4.5	(c1)	4 0	(c2)	4.3	
8. BOLTING: (Give First Water Date: 05/24/2005) NOTE: First Water Date is the date seed first receives adequate moisture to germinate. This can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1) 96 (c1) 93 (c2) 91 BOLTING CLASS: 1 = Very Slow 3 = Medium 2 = Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	CORE HEIGHT FROM BASE	OF HEAD TO APEX:						
Can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1)		(a1)	3 9 mm	(c1)	4 2 mm	(c2)	37 mm	
Can and often does equal the planting date. NUMBER OF DAYS FROM FIRST WATER DATE TO SEED STALK EMERGENCE: (summer conditions) (a1)	8 BOLTING: (Give First Water De	ote: 05/24/200.	S MOTE 5	First Mistor Date :	s the data as ad E	et roooluse ==	Nograta maintur-	to corminate This
(a1) 96 (c1) 93 (c2) 91 BOLTING CLASS: 1 = Very Slow 3 = Medium 5 = Very Rapid 2 = Slow 4 = Rapid (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	Clive I list vvalid! Da			can and often do	es equal the plant	ing date.	reduare moistale	to germanate. This
BOLTING CLASS: 1 = Very Slow 3 = Medium 5 = Very Rapid 2 = Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	NUMBER OF DAYS FROM FIRS	ST WATER DATE TO	SEED STALK EM	ERGENCE: (sum				
2 = Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:		(a1)	96	(c1)	93	(c2)	91	
2 = Slow 4 = Rapid (a1) 2 (c1) 2 (c2) 3 HEIGHT OF MATURE SEED STALK:	BOLTING CLASS:	1 = Very Slow	3 = Medium	5 = Very Ra	apid			
HEIGHT OF MATURE SEED STALK:		2 = Slow				•		
1/4 /2/ 97		(a1)	2	(c1)	[之]	(c2)	[3]	
	HEIGHT OF MATURE SEED STA	ALK:						. (
	ST-470-1 (04-05) designed by the Plant Variet	ly Protection Office using f			121		97	Page 4 of 9

(a1)

cm

(c1)

8. BOLTING: (continued)

SPREAD OF BOLTER PLANT: (At widest point)

(a1)

2 2 cm

18 cm

BOLTER LEAVES:

1 = Straight

2 = Curved

MARGIN:

1 = Entire 2 = Dentate

COLOR:

1 = Light Green

2 = Medium Green 3 = Dark Green

2 (a1)

BOLTER HABIT:

TERMINAL INFLORESCENCE:

1 = Absent

2 = Present

2 (a1)

(c1)

2 (c2)

LATERAL SHOOTS:

1 = Absent

2 = Present

2 (a1)

(c1)

BASAL SIDE SHOOTS:

1 = Absent

2 = Present

(c1)

(c2)

9. MATURITY: (earliness of harvest-mature head formation)

NOTE: Complete this section for at least one season.

SEASON	А	PPLICAT			MOS	ST SIMILAR V		STANDARD F	REGIONAL CHECK V	
		No.	of Days	'		No. of Day	's'		No. of Days	1
Spring	A gg	3 8	71	c 71	A 96	⁸ 70	c 69	A 96	8 70	c
Summer	A 74	В	69	c 67	A 74	B 69	c 67	A 74	8 —	c _
Fall	450		-	_	-	-	_	_	_	
Winter	A 126		_	-	A 121	-		A 126	-	_

First Water Date to Harvest

Give Planting Date(s) and Location(s):

Spring:

A: 2/13/06 (SALINAS, CA), B: 4/21/05 (SANTA MARIA, CA), C: 4/14/04 (SALINAS, CA)

Summer:

A: 6/18/05 (SAN JUAN BAUTISTA) B: 5/20/05 (SALINAS, CA), C: 6/1/04 (SALINAS, CA)

Fall: Winter:

A: 12/16/05 (SAN JUAN BAUTISTA, CA)

10. ADAPTATION:

PRIMARY REGIONS OF ADAPTATION (tested and proven adapted):

0 = Not Tested

1 = Not Adapted 2 = Adapted

Southwest (CA and/or AZ desert)

West Coast

Northeast

North Central

Southeast

Other (Specify)

10.	. ADAPTATION: (Continued)				
SE	ASON: CALINAS	SANTA MARIA,			
	2 Spring (Area HURON,)	'UMA)	2 Fall (Area SALI	NAS, SANTA MARI	A)
	2 Summer (Area SALINAS, SA	NOTA MARIA)	Winter (Area	_	
	· · · · · · · · · · · · · · · · · · ·				
-	GREENHOUSE: 0 = Not	Tested 1 = No	t Adapted 2 = Ada	apted	
	SOIL TYPE: 1 = Min	eral 2 = Org	ganic 3 = Bot	th	
11.	VIRAL DISEASES:		· more and		
	1 = Immune 3 = Resistant	5 = Moderately Resist	ant/Moderately Susceptible	7 = Susceptible	9 = Highly Susceptible
	Big Vein	(a1) 7	(c1) 7	(c2) 7	0=not tested
	Lettuce Mosaic	(a1) 7	(c1) 7	(c2) 7	
	Cucumber Mosaic	(a1) O	(c1) O	(c2) O	
-	Tomato Bushy Stunt, cause of dieback	(a1) 0	(c1) O	(c2) O	
	Turnip Mosaic	(a1) 0	(c1) 0	(c2) O	
	Beet Western Yellows	(a1) 0	(c1) O	(c2) O	
	Lettuce Infectious Yellows	(a1) 0	(c1) O	(c2) O	
	Other (Specify)	(a1) (0	(c1) O	(c2) O	
	EUNCAL (PACTERIAL DISEASES)				
12.	FUNGAL/BACTERIAL DISEASES: 1 = Immune	5 - Madarataly Basists	ent/Madarataly Cycanomible	7 - Oursestible	O – I lizabili. Ovacantibla
	Corloy Poot Pot		ant/Moderately Susceptible		9 = Highly Susceptible
	(Races: CA I	(a1) 3	(c1) 3	(c2) 3	0 = not tested
AD	Downy Mildew	VI (a1) 3	(c1) 3	(c2) 7	
107	(Races: CA I and II	(a1) [3]	(c1) [3]	(c2) [+]	
	Powdery Mildew	(a1) O	(c1) O	(c2) O	
	Sclerotinia Drop	(a1) 7	(c1) <u>7</u>	(c2) 7	
	Bacterial Soft Rot (Pseudomonas spp. and others)	(a1) O	(c1) <i>O</i>	(c2) <i>O</i>	
	Botrytis (Grey Mold)	(a1) 0	(c1) O	(c2) 0	
	Verticillium Wilt	(a1) 0		` ' 🗖	
	Bacterial Leaf Spot	(a1) 0	(c1) <i>O</i>		
	Anthracnose	(a1) O	(c1) <i>O</i>		
	Other (Specify)	(a1) O	(c1) O	(c2) [<i>O</i>]	
40	-	(61) [2]	(61)	(62)	·
13.	INSECTS:	f — hiladayatalı. Davist	-484-d4-b- O	7 O	N. Illahla Ossassatilla
	1 = Immune 3 = Resistant	o = Moderately Resistal	nt/Moderately Susceptible	7 = Susceptible \$	= Highly Susceptible
	Cabbage Loopers	(a1) O	(c1) 7	(c2) O	0 = not tested
	Root Aphids	(a1) 0	(c1) O	(c2) O	
	Green Peach Aphid	(a1) 7	(c1) 7	(c2) 7	
	Lettuce Aphid	(a1) 7	(C1) 7	(c2) [7]	11 .

	Pea Leafminer Other (Specify)	(a1) O (a1) O	(c1) 0 (c1) 0	(c2) 0 (c2) 0	
14.	PHYSIOLOGICAL STRESSES:				
	1 = Immune 3 = Resistant	5 = Moderately Resistant/	Moderately Susceptible	7 = Susceptible 9 = H	lighly Susceptible
	Tipbum	(a1) 5	(c1) 5	(c2) 5	0 = not tested
	Heat	(a1) O	(c1) O	(c2)	
	Drought	(a1) 0	(c1) O	(c2) 0	
	Cold	(a1) O	(c1) O	(c2) O	
	Salt	(a1) O	(c1) 0	(c2) O	
	Brown Rib (Rib Discoloration, Rib Blight)	(a1) <u>5</u>	(c1) [5]	(c2) <u>5</u>	
	Other (Specify)	(a1) <u>(</u>	(c1) Ø	(c2) O	
15.	POST HARVEST STRESS:				
	1 = Immune 3 = Resistant	5 = Moderately Resistant/M	loderately Susceptible	7 = Susceptible 9 = H	ighly Susceptible
	Pink Rib	(a1) <u>5</u>	(c1) <u>5</u>	(c2) <u>7</u>	0 = not tested
	Russet Spotting	(a1) <u>5</u>	(c1) 5	(c2) 7	
	Rusty Brown Discoloration	(a1) 5	(c1) 5	(c2) O	
	Internal Rib Necrosis (Blackheart, Grey Rib, Grey Streak)	(a1) 5	(c1) 5	(c2) O	
	Brown Stain	(a1) 5	(c1) 5	(c2) O	
		•			

16. BIOCHEMICAL OR ELECTROPHORETIC MARKERS:

17. COMMENTS:

SUGGESTED CHECK VARIETIES

TYPE

Cutting/Leaf

Butternead 2

3 Bibb

4 5 Cos or Romain

Great Lakes Group

Vanguard Group

7 Salinas Group

Eastern Group 8

9 Stem

10 Latin CHECK VARIETY Waldmann's Green

Dark Green Boston

Bibb

Parris Island

Great Lakes 659-700

Vanguard

Salinas

Ithaca

Celtuce

Little Gem

REFERENCES

Bowring, J.D.C., 1969, "The Identification of Varieties of Lettuce (Lactuca Sativa L.)". Journal of the National Institute of Agricultural Botany 11:499-520. National Institute of Agricultural Botany, Cambridge, UK.

Davis, R.M., K.V. Subbarao, R.N. Raid, and E.A. Kurtz, 1997. "Compendium of Lettuce Diseases". APS Press, St. Paul, MN.

Michelmore, R.W., J. M. Norwood, D.S. Ingram, I.R. Crute and P. Nicholson. 1984. "The interitance of virulence in Bremia lactucae to match resistance factors 3, 4, 5, 6, 8, 9, 10, and 11 in lettuce (Lactuca sativa)", Plant Pathology 32:176-177.

Norwood, J.M., R.W. Michelmore, I.R. Crute and D.S. Ingram. 1983. "The inheritance of specific virulence of Bremia lactucae (Downy Mildew) to match R-factors 1, 2, 4, 6, and 11 in lettuce (Lactuca sativa)". Plant Pathology 32:176-177.

Rodenburg, C.M., et al., 1960. "Varieties of Lettuce. An International Monograph", Instituut voor de Verdeling van Tuinbouwgewassen (IVT), Wageningen, NL.

Ryder, E.J., 1999, Lettuce, Endive, and Chicory, CABI Publications, Wallingford, UK.

Exhibit C - Tellmark

Fourth leaf from 20 day-old seedlings of 'Tellmark'





Exhibit C - Tellmark

Frame leaves of 'Tellmark'







Exhibit C - Tellmark

Heads of 'Tellmark'





16

Exhibit D - Tellmark

Table 2a: Bremia lactucae Screening- Isolate CA-V

	Rep1		Rep 2	2	Rep 3	3]		
Cultivar	+	-	+	- ,	+	-	# Resistant	# susceptible	% Resistant
Tellmark-foundation seed	0	16	0	16	0	16	48	0	100%
Tellmark-production seed 2004	0	16	0	16	0	16	48	0	100%
Tellmark-production seed 2005	0	16	0	16	0	16	48	0	100%
Telluride	0	16	0	16	0	16	48	0	100%
Hallmark	16	0	16	0	16	0	0	48	0%

Table 2b: Bremia lactucae Screening- Isolate CA-VI

	Rep1		Rep 2	2	Rep :	3			·
Cultivar	+	-	+	-	+	-	# Resistant	# susceptible	% Resistant
Tellmark-foundation seed	0	24	0	24	0	24	72	0	100%
Tellmark-production seed 2004	0	24	0	24	0	24	72	0	100%
Tellmark-production seed 2005	0	24	0	24	0	24	72	0	100%
Telluride	0	24	0	24	0	24	72	0	100%
Hallmark	24	0	24	0	24	0	0	72	0%

Table 3: Molecular Marker for Bremia Resistance in Telluride

Cultivar	Y/N*
Tellmark-foundation seed	Υ
Tellmark-production seed 2004	Υ
Tellmark-production seed 2005	Υ
Telluride	Υ
Halimark	N

Y= marker present N= marker absent

Table 4: Corky Root Bioassay- Sphingomonas suberifaciens-Isolate 'CA-I'

	Rep1		Rep 2	2	Rep :	3	1		
Cultivar	+	-	+	-	+	-	# Resistant	# susceptible	% Resistant
Tellmark-foundation seed	0	16	0	16	0	16	48	0	100%
Tellmark-production seed 2004	0	16	0	16	0	16	48	. 0	100%
Tellmark-production seed 2005	0	16	0	16	0	16	48	0	100%
Telluride	0	16	0	16	0	16	48	0	100%
Hallmark	e 0	16	0	16	0	16	48	. 0	100%

Table 5: Molecular Marker for cor gene

Cultivar	Y/N*
Tellmark-foundation seed	Υ
Tellmark-production seed 2004	Υ
Tellmark-production seed 2005	Υ
Telluride	Υ
Hallmark	Υ

Y= marker present N= marker absent

Seedling measurements at 4th leaf stage

ratio (I/w*10) 12.00 13.75 13.93 16.25 16.25 17.74 17.50 17.50 11.76 11.76 11.76 11.76 11.76 width (cm) 25.54 25.54 25.54 25.55 length (cm) 4.13 3.81 3.81 3.81 3.81 3.49 4.45 3.49 4.45 4.13 3.18 3.49 3.81 3.81 ratio (I/w*10) 14.44 15.00 17.10 15.56 15.50 17.50 width (cm) 2.554 2.286 2.286 2.296 2.254 3.02 2.254 2.254 2.254 2.254 3.302 3.302 2.366 3.362 3 length (cm) 4,45 4,45 3,81 3.97 3.81 4.45 3.81 3.97 3.81 4.45 ratio (I/w*10) 12.08 12.78 13.00 15.24 14.00 11.50 12.00 17.50 14.38 12.73 12.00 13.33 9.59 12.09 12.73 12.00 10.77 13.81 Tellmark width (cm) 3.81 2.86 3.18 3.33 3.18 3.18 2.54 2.54 3.49 3.18 3.34 3.49 3.18 3.81 length (cm) 4.45 4.60 3.65 4.13 5.08 3.65 4.13 4.45 3.81 3.65 4.45 4.45 3.65 4.61 3.81 Plant # Mean

<u>- Tellmark</u>

Statistical Comparison Tellmark vs. Telluride (table 1)

E (crit) 05 = 4.40 E (crit) 04 = 4.5F	n = 20	E (crit) 100 - 4:10; F (crit) 101 - 7:55	F (crit) $.05 = 4.10$, F (crit) $.01 = 7.35$
n = 20			n = 20
Cut date- 6/30/05		Cut Date- 4/21/06	Cut Date- 5/19/06
Sow date- 4/21/05	Sow date- 6/8/05?	CO	
SMA1 Santa Maria, CA	SJB2 San Juan Bautista, CA	SJB3 San Juan Bautista, CA	SAL4 Salinas, CA
SMA1	SJB2	SJB3	SAL4

Trial		SMA1	S	SJB2	S	SJB3	A.S.	SAI 4
Trait	Tellmark	Telluride	Tellmark	Telluride	Tellmark	Telluride	Tellmark	Telluride
Spread of Frame Leaves :						2	100	מומווספ
Mean (cm)	57.0	54.5	62.6	51.7	52.2	49.0	53.9	50.3
Std. Dev.		2.1	3.1	2.4	60	er.	3.7	X
ANOVA, F (calc)	8.48 **	*	154.59 **		8.72 **		15.11 **	
Head Weight:	-							
Mean (grams)	846.9	793.4	959.7	779.1	522.7	514.4	5283	5703
Std. Dev.	161.7	109.9	147.4	130.8	85.4	096	109.4	95.2
ANOVA, F (calc)	1.50 ns	us	16.81	**	0.08 ns		1.67 ns	
Head Diameter:								
Mean (cm)	15.0	13.3	15.9	14.1	12.9	11.8	13.4	12.0
Std. Dev.	0.8	1:1	0.8	1.0	<u>.</u>	, c	ξ α 2	2 6
ANOVA, F (calc)	28.34 **	**	37.31 **		10.79 **		1.03 ns	
Head Height:								
Mean (cm)	14.0	14.0	16.7	15.1	12.1	11.2	11.7	7 2 2
Std. Dev.	6:0	9.0	9.0	0.	0.9	8.0	80	20
ANOVA, F (calc)	0.01 ns	ns Su	29.60		.95	*	24	*
Core Diameter :								
Mean (mm)	32.0	35.2	36.7	33.5	30.6	30.3	27.7	30.5
Std. Dev.	4.3	2.0	2.3	2.0	2.5	2.5	3.7	2.6
ANOVA, F (calc)	9.14	* *	21.78 **		0.10 ns		.27	r **
Core Length :								
Mean (mm)	38.4	46.9	67.8	62.1	25.2	28.3	23.8	31.0
Std. Dev.	9.5	7.6	9.5	13.5	4.4	6	4.4	7.2
ANOVA, F (calc)	10.28 **	**	2.35 ns	ns Su	5.76	*	4	!
Leaf Color:								
RHS Color Standard	137C	138A	137C	138A	137C	138A	137C	138A

^(*) statistical significance at probability level of 0.05 (**) statistical significance at probability level of 0.01 (ns) statistical not significant

¹⁹

- Tellmark

Data Set Plant Charachteristics

Evaluation date- 6/30/05	evaluation date- 8/22/05	valuation date- 4/21/06	valuation date- 5/19/06
Sow date- 4/21/05 Eval		Sow date- 12/16/06 Evall	ш
Santa Maria, CA	San Juan Bautista, CA	San Juan Bautista, CA	Salinas, CA
•••	•••		••
SMA1	SJB2	SJB3	SAL4

_																							Æ17	n Hj	18	Ψ,	F' 7	¥ 1	ij	20	őkas	. F.,	ÿ	
	-	Hallmark		٠																		143A	5											
(31.0)	יוסוי (וארויט	Felluride																				138A	}											
	<u>-</u> ادّ	23.1 35 30.7 22.4 52.6 60.4																				137C)					•						
100		allmark	- 60.	1.0.4 2.0.4 8.0.4	24.0	1 %	3 8	47.7	47.1	64	32.6	36.8	32.9	56.1	414	24.7	38.9	42.7	25.0	48.2	37.9	420	8.7	37.1	54.3	71.4	52.5	50.6	34.5	41.4	72.3	50.4	37.8	43.1
Core Length (mm)		elluride	33.0	1 6 1 0	1 7 7	5.4		41.5	38.5	43.9	41.5	39.9	8.64	57.2	45.7	7 07	47.8	37.8	57.3	52.8	30.0	46.9	7.6	42.8	38.7	60.1	63.3	65.5	69.4	51.3	8.69	54.6	47.4	48.2
200		22.4	4.20 20 0	23.4	42	40.5	20.7	43.2	42.1	32.4	50.8	39.8	38.2	38.7	42.3	53.2	35	51.4	42.2	44.6	35.2	38.4	92	80.1	61.7	72.5	82.3	88.5	76.8	6.69	57.7	56.8	66.8	68.7
(m	1111	30 7		33.8	75	26.2	33.7	92	29.8	35.8	33.1	31.4	31.4	34.2	37.3	30.6	32.6	32.5	33.9	35.2	36.6	32.8	2.8	36.8	41.2	39.1	31.7	39.3	30.7	37.2	42.1	37.3	31.5	38.2
Core Diameter (mm)	J opinil	35	34.9	34.3	31.3	37.2	37.2	35.4	33.4	35.7	37	38.1	35.2	37.6	38.3	34.1	35.2	32	35.1	32.6	34.8	35.2	2.0	32.3	36.1	30	37.4	32.4	32.7	35.2	32.5	30.7	33.5	34.5
Core	Ilmort T	23.1	30.1	24	29.1	33.6	29.3	37.3	34	26.2	39.9	31.2	35.3	35.2	37.1			29.4	33.4			32.0	4.3	34.2	36.3	37	35.4	40.2	38.7	37.2	36.8	14	34.2	38.4
-			. 6	2.2	5	13.2	13.7	13.4	14.1	13.4	14.3	14.1	12.3	12.9	13.6	13.4	13.7	13.6	13.1		13.5	13.5	9.0	16.1	15.2	15.3	14.7	15.9	12.8	1.4	15.3	8.41	13.1	5.3
Head Height (cm)	Inride Ha	13.4	6.4	4	13.4	14.5	5.5	•	13	13.7	16.4	14.3	14.4	13.9	13.8	14.4	13	12.9	`		13.3	14.0	9.0	14.4	14.6 1	14.6 1	17 1	14.8	15.1	5.1	16 1	13.3	14.7	14.8
Head H	Hallmark Telluride Hallmark	12.4	-		·	14.1	14	5.8	14.1	13.2	14.3	14.2	14.1	13.5	14	14.3	13.4	14.5	13.2	15.2	15.4	14.0 1	0.9	16.1		17.4 1,	15.7	16.8	16.4	17 1	6.5	17.2 1	16 1	17.7
L	ark Tell	9								-												\vdash					_		<u>`</u>		_	_		_
ster (cm)					12.8	13.1	13.6	14.4	15	15.2	13.9	14.2	12.4	13.9	14.2	12.6	14.9	13.4	13	13.7	12.8	13.7	0.8	16.4	17		15.2	14.9	14.4	15.2	15.4	14.8	13.8	14.2
Head Diameter (cm)	rk Telluri	13	13.1	12.4	12.2	14.5	5	14.7	12.5	13	15	12.3	13.9	12	15.7	14.9	13.5	13.4	12	13.7	12.1	13.3	1.1	14.4	14.1	14.6	12.8	14.1	16	13	14.2	12.6	15.1	12.6
H	k Tellma	14.8	16	13.7	14.9	15.5	15.2	14.6	15		14.1	14.9	13.5	16.3	15.9	16.1	15.4	14.5	15.4	16.2	14.3	15.0	0.8	13.9	16.4		15.8	14.7	15.2	16.9	91	14.9	16	16.9
grams)	Hallmai	698.6	732.8	722.9	656	632.6	697.1	897.5	722.7	869.2	687	671.3	765	799.8	843.6	734.3	729.8	713.4	645.7	636.9	891.5	737.4	82.7	1083.9	1253.6	1065.3	538.6	1283.5	637.8	815	950	1106.3	537.2	1029.4
Head Weight (grams)	Telluride	744	721.9	767.9	641	807.2	893.2	988.9	598.3	825.5	1020.4	855.1	789.6	857.9	906.8	771.3	773.4	606.9	768.9	774.5	754.5	793.4	109.9	676.5	926	552.4	997.5	726.9	888.9	818	933.8	621.5	762	688.7
Head	Tellmark	787.1	584	690.3	882.8	746.9	722.4	1001.1	876.4	646.8	1126.5	624.1	874.8	880	1226.5	881.5	794.2	822	845.7	1003.9	910.7	846.9	161.7	778.9	930	1200	1022.7	1048.1	764.9	1172.1	752.7	1040	963.1	736.3
aves (cm)	Hallmark	51.9	54.3	20.2	52.2	55.5	54.5	55.1	20.2	55.3	54.4	58	51.3	51.5	52.1	54.6	50.9	58.5	53.7	53.4	53.1	53,6	2.3	49.9	60.7	28	56.8	6.09	46.8	53.8	58.9	54.5	54.2	23
Spread of Frame Leaves (cm)	Telimark Telluride Hallmark Tellmark Telluride Hallmark Telluride	52.3	52.9	50.1	55.7	55.2	54.4	9.99	56.5	54.2	23	½	54.3	56.3	53.9	9	55.6	51.4	55.6	54.7	53.8	54.5	2.1	48.5	54.8	51.9	51	49.7	53.8	50.5	58.1	50.8	53.2	50.3
Spread o	Tellmark	6.53	51.3	56.3	54.2	57.8	61.4	58.4	54.7	58.9	26	53.4	54.8	59.3	62.3	57.4	58.7	52	58.3	8.09	59.9	57.0	3.1	55.3	64.5	65.2	62.8	61.9	55.1	58.9	63.3	62.7	65.6	65.7
推 性		1	7	ო	4	ις.	ဖ	7	ω	0	6	Ξ	5	<u>რ</u>	4	ਨ	16	17	8	9	ឧ		1	_	7	ი .	4	2	ဖ	7	œ	တ	0	=
Trial#		SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	SMA1	Mean	StDev	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2

20

- Tellmark

Data Set Plant Charachteristics (continue)

г	_	रा									<u> </u>		_										_			É) (6	0	0	87) š) (3			
		165 149 137 366 237 375 C										143A																			-	-	1434	5					
į	Color (KHS)	elluride									, ,	₹85.																					1384	j					
Č	3 - 	mark									ç	<u>ع</u> رد																					137C						
-	+	mark	58.2	<u> </u>	50.5	, i	7.64	34.7	72.4	37.6	20.2		200	36.3	34.2	29.9	35.4	: 4	. "	, ,	, c	23.8	. ~		: 4		rن 	 		4	· c		╀		 	~	· •		ئة —
(h)		ide Hall									ı																		36.3				ı		``				3 27.5
1			4. 4. c		0.50	9 8	20.00	7.00	20.3	62.4	92.6	13.5	2 5	27.8	32.4	29.5	30.5	36.4	34.3	2 8 8 8	33.4	24.9	30.8	24.7	23.6	24.5	34.3	24 7	23.2	27.5	25.4	26.2	283	3.9	32.4	32.1	29.2	36.3	30.8
	١	e g	00 4 4.00 0	40.0	7.10	67.5	5.70	70.4	3 6.6	72.0	23.3) r	28.4	26.7	20.1	32.4	22.7	28.6	29.1	0 6	16.8	24.7	22.6	24.4	30.2	23.1	32.6	20.1	26.5	21.6	25.1	27.8	25.2	4.4	20.0	26.3	27.5	26.2	25.4
(mm)	, ,	nailmar	5. 5. C. 15.	- o	30.00 80.00	9 6	0 0	32.0	0.9.0	, ç	36.7	3.7	30.1	25.5	28.3	27.8	33.1	31.1	33.8	29.7	30.1	29.7	31.2	26.3	27.1	27.3	32.4	31.7	37.2	28.2	29.3	29.2	30.0	2.8	23.2	24.8	24.0	12.7	28.3
Core Diameter (mm)	off, side		32.3	34.5	- 8	2 4 6	2 6	5. 5. 5. 5. 5. 5.	2. t. 6.	0 16	3 2	2.0	26.2	33.4	32.1	31.2	34.1	32.2	30.7	30.2	32.1	엁	29.6	29.7	28.3	25.9	30.2	33.6	28.2	29.6	26.8	28.4	30.3	2.5	31.0	29.9	30.0	35.1	31.2
Core D	T viscoul	26 F	34.2	34.1	38.5	37.2	2 7 5 2 7 2	3.7.5	27.6	30.3	36.7	2.3	29.2	31	26.2	29.3	34.2	30.5	32.3	28.4	26.8	29.5	30.3	34.5	34.7	29.6	29.4	30	31.2	28.2	32	34.2	30.6	2.5	27.0	28.1	31.4	30.0	29.8
-	mork	13.7									╀		\vdash		12.6		13.2	13	12.7			12.7	12.5	11.4	11.5	12.4	11.9	13.5	10.9	13.1	12.1	12.7	12.6	0.8	<u> </u>	10.8	11.4		11.9 2
Head Height (cm)	ride Hall	9	•	·	,	•		•	·																	•	•			•	·								
lead Hei	ırkİTellı	14 9			-	,-	•			_			12.1	12.7	11.8	11.7	T	11.5	12	11.4	•	10	11.5	6.6	10.5	9.6	11.7	10.8	11.3	10.5	10.9	11.9	11.2	0.8	11.0	10.9	11.0	11.8	10.0
	k Tellms	16.5	17.3	16.4	16.8	18.4	15.9	15.5	16.1	17.7	16.7	0.8	12.8	1.1	12.7	12	12.7	13.8	12	9.7	11.2	12.5	11.9	12.9	11.9	11.6	12.2	12.5	12.4	10.7	12.6	12.1	12.1	0.9	12.0	13.1	12.4	12.5	12.7
(Gm ²)	Hallmar	13.9	15.5	14.7	13.9	14.1	14.2	17.1	15.9	4	15.0	1.0	11.2	12.9	11.2	10.7	12.2	12.1	11.8	12.9	5	12.4	10.7	12.1	12.6	10.7	10.7	12	10.1	11.4	11.9	13.2	11.8	6.0	11.6	13.7	12.5	13.3	13.2
Head Diameter (cm)	Felluride	15.3	13.5	15	14.4	13.4	15.9	13.5	13.7	14.2	1.4	1.0	12.2	<u>ლ</u>	75	13.2	13.3	12.7	6.6	11.9	12.7	10.9	11.6	10.3	9	;	1.5	11.4	12.2	11.7	11.6	12	11.8	1.0	13.9	12.0	13.8	14.0	13.0
Head	ellmark 1	16.4	16.2	16	17	16.5	15	17	15.5	15.9	15.9	9.0	13.2	7	10.8	11.2	15	14.4	13.8	12.5	13.9	12	12.9	12.8	12.7	13.8	13.9	12	12.5	13.5	12.8	12.4	12.9	1.1	13.4	14.0	14.2	14.0	13.5
(SI	Imark Te	727.3	902.1	657.1	817.1	893.1	661.2	1005	782.6	754	875.0	220.0	575.7	530.9	561.5	518.1	261	646.5		619.4								<u></u>				612.9	548.7	88.7	480.0				666.7 1
Head Weight (grams)	ride Hal	5.6 7.	705.6 90																																-		-		
ad Weig	ark Tellu	2 845.6		.2 795.3	7 779.9	.5 782.9	.9 826.3	.8 718.2	2 539.7	4 995.5	7 779.1	4 130.8								5 591.1						.,	533	462.4				487.5	514.4	96.0					623.0
윈	rk Tellm	856.2	818.7	1038.2	1042.7	1015.5	1007.9	1165.8	792.2	1048.4	959.7	147.4	524.5	471.1	456.5	421.7	743.5	641.2	624.5	403.5	399.4	521.1	541.3	523	641.1	520.8	488	489	202	482.1	523.4	533.5	522.7	85.4	558.0	669.2	684.0	613.0	676.0
Spread of Frame Leaves (cm)	Tellmark Telluride Hallmark Tellmark Telluride Hallmark Tellmark Telluride Hallmark	51.5	58.7	55.6	56.5	53.4	55.3	57.2	48.9	48	54.6	1.4	51.5	46.9	49.9	49.7	47.8	50.2	54.3	54.1	52.5	53.3	4 4.8	47.5	48	51.5	52.8	55.7	44 .3	48.8	46.9	20	50.0	3.2	45.0	54.3	50.2	53.9	51.0
Frame Le	Telluride	25	48.7	49.9	51.7	49.8	52.1	52.5	49	55.2	51.7	2.4	20.8	53.4	51.8 8. i	51.3	48.9	48.2	46.5	50.8	54.2	45.3	49.8	43.9	46.9	40.8	48.8	46.9	51.5	52.8	49.4	48.7	49.0	3.3	48.6	49.2	52.4	49.8	20.7
Spread of	ellmark	60.5	64.5	64.9	62.8	64.7	61.8	63.8	62.5	65.3	62.6	3.1	54.8	51	47.4	25	51.8	56.8	49.9	49.5	49.3	53.2	50.5	57.5	54.6	ဥ	53.9	<u>,</u>	26	53.8	4 4.8	49.2	52.2	3.3	57.0	51.5	56.8	62.3	ე <u>დ</u> . ე
# _H	7		5	4	ئ	9	17	8	9	20		+	- 7	N (· ·													1 0				8		\dashv					
Tria#		SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	SJB2	Mean	Stdev	SJB3	200	2000	SALS GILO	2000	2003	SJB3	SJB3	SB3	SJB3	2002	SJB3	SJB3	V.JB3	2000	2000	200	SJB3	SJB3	SJB3	Mean	Stdev	SAL4	SAL4	SAL4	SAL4	OAL4

- Tellmark

Data Set Plant Charachteristics (continue)

	(S)	Tellmark Telluride Hallmark																	1	143A			
	Color (RHS	Tellurid																	. 007	138A			•
		Tellmark																	4070	2/51			
	mm m	Hallmark	22.7	31.4		0.02	28.5	29.4	9.87	24.7	20.8	26.1	22.4	22.6	2 2	2 4.0	0.77	4 L	4.72	4.0.	3.8		
- Canada	Core Length (mm)	Telluride	32.4	27.8	0 0) L	20.0	27.4	707	λ. 	5.4°	35.7	19.9	26.5	27.0	1 6	27.0	5.75	27.00	<u>, </u>	7.	,	
3	3	Tellmark	31.2	24.4	110	. c	7.5.	7.07	20.0	7.5.d	2.23	21.4	20.7	33.2	23.1	28.2	20.6	23.7	23.0	0.54	‡	8	
(mm)		Talling Lellingrik Fellunde Hallmark Fellunde Hallmark Tellunde Hallmark Tellunde Hallmark	19.5	30.2	30 3	2 40	4. 0	7.0.7	20.0	7 0 7	7.07	23. 20.	30.1	28.3	26.3	280	1 0	. 00	25.4	† ,	,	, , ,	
Core Diameter (mm)	ומוומנו	Telluride	31.1	29.2	34.1	300	20.00	20.2	24.7		0.20	0.40	25.6	29.1	30.0	29.1	30.5	200	30.5	3 6	1	, , ,	
Core	3	ellmark	29.9	30.1	18.7	75.6	24.0	5 2	27.2	1 α	7 0	0.12	22.8	35.0	24.9	33.4	25.8	28.2	27.7			1 10	
(mg		Tallmark	10.8	13.0	11.2	. 2	5 0		12.6	2 4	2 4	2 :	 	10.9	10.5	1,1	10.2	1.3	114	σ C		43.0	
Head Height (cm)		elluride	10.5	12.3	11.9	10.0	10.6	10.7	1 1	, C	5 5) (- ;	7.1	12.2	12.1	10.8	11.7	1.5	112	20		12.0	
Head	7	CIIIIBIL	11.7	11.2	8.6	11.4	10.9	10.5	1.5	12.2	100	1 6	3.0	12.0	10.9	12.0	11.2	11.4	11.7	8.0		13.6	
(cm)	- Form	iall lial k	12.1	14.1	14,4	14.7	12.1	13.4	12.8	14.2	146		<u>4</u>	12.8	14.1	13.3	11.6	14.5	13.4	0.0		13.5	
_			12.5	13.0	13.0	13.5	13.6	13.4	13.2	14.4	12.5	5	7.7	13.8	13.0	12.5	13.2	13.0	13.2	0.7		13.1	
Head	elimark	A CONTRACTOR	13.3	13.7	12.9	13.2	13.4	12.0	12.3	12.5	13.2) !	14.0	13.7	15.5	12.5	12.7	13.4	0.8		14.3	
(Sur	lallmark T	0.00	0.000	753.1	586.6	580.0	476.6	708.0	655.0	839.0	688.1	7107		554.3	638.8	693.2	391.6	0.069	602.3	126.3		8.069	
Head Weight (grams)	elluride	1007	400.0	504.0	588.5	602.1	505.5	484.5	572.8	741.2					470.5	494.2	508.0	430.8	570.3	95.2		664.3	
Head W	ellmark	516.0	0.0	515.8	329.9	422.4	441.8	410.0	440.0	447.1	638.0				513.5	658.0	409.8	495.6	528.3	109.4		714.4	
(es (cm)	Tellmark Telluride Hallmark Tellmark Telluride Hallmark Tellmark Telluride	23 B	2 6	χ. Σ.	48.5	48.2	52.7	47.4	52.9	54.8	52.6	47.4			51.4	51.3	52.5	47.8	51.1	2.8		52.3	
rame Leav	elluride	50.0	1 9	7.04	8.6	52.9	54.8	49.5	48.7	48.6	49.3	48.5	70.0	P (20.8	50.5	52.4	51.9	50.3	1.8		51.4	
Spread of Frame Leaves (cm)	ellmark	514		7.00.	54. 8	52.0	53.8	54.5	51.2	51.0	47.8	54.9	78.0	5 1	55.6	51.5	49.7	48.8	53.9	3.7		56.4	
±	_	g) h	- (xo.	თ	9	Ξ	12	13	4	15	9	2 [1	8	19	20				ot ot	
Llat		SAL4	7 70	ָרְיָּרְיִּרְיִּרְיִּרְיִּרְיִּרְיִּרְיִּרְיִ	2AL4	SAL4	SAL4	SAL4	SAL4	SAL4	SAL4	SAL4	SAIA	1	SAL4	SAL4	SAL4	SAL4	Mean	StDev		Mean tot.	

- Tellmark

Data Bolting Charachteristics

Location :

San Juan Bautista, CA

တိ

Sow Date: 5/24/2005

些	# Days to	# Days to Seed Stalk Emergence	пегделсе	Height of	Height of Mature Seed Stalk (cm)	Stalk (cm)	Spread	Spread of Bolter Plant (cm)	it (cm)
	Tellmark	Telluride	Hallmark	Tellmark	Telluride	Hallmark	Teilmark	Telluride	Hallmark
_	86	84	9	114.96	100.32	89.28	24.48	23.52	15.6
7	100	87	86	107.52	113.52	108.96	30.48	21.36	18.96
<u>ო</u> .	86	101	86	106.08	126.72	75.36	24.72	22.32	12.48
4	88	100	94	104.4	129.36	72.72	23.76	22.8	10.56
, O	86	91	91	106.32	115.68	89.28	17.52	19.44	16.32
ဖ	86	8	91	104.4	115.92	79.92	23.04	22.08	18 72
7	100	06	9	96.72	118.8	94.8	23.28	20.64	18.72
œ	86	06	<u>ه</u>	123.6	126.72	98.88	21.36	21.6	16.08
တ	98	06	86	114	121.92	106.32	23.52	24.48	16.32
9	86	86	<u>Б</u>	113.52	111.36	89.28	19.92	19.92	15.36
7	91	06	88	119.76	119.52	82.32	20.16	22.08	16.56
72	9	94	86	123.12	123.12	98.86	19.68	19.68	16.32
<u></u>	86	06	86	110.88	128.4	107.76	23.52	21.12	€
4	88	86	ক	115.2	128.16	ı	17.28	20.16	٠ ؛
5	88	91	98	123.6	118.8	95.04	20.64	19.92	16.56
9	26	108	86	113.28	132.48	96.72	20.16	20.4	19.68
7	86	86	8	114.24	130.56	100.8	22.56	23.52	23.76
<u>&</u>	88	100	98	119.76	113.04	94.08	24.72	22.08	19 44
6	86	91	91	125.04	128.16	88.32	24.48	22.8	16.8
20	87	æ	8	120	115.2	110.64	22.08	21.84	18.24
77	100	91	စ္ထ	119.04	118.8	121.92	24.72	20.64	21.36
22	86	•	\$	111.6	1	109.2	24.96	•	24.48
23	86	1	91	112.32	•	115.68	27.12		21.6
77	88	ı	90	113.52	•	114.24	19.44	1	17.52
Mean	95.7	92.7	91.2	113.9	120.8	97.4	22.7	21.5	17.8
StDev	43	0	1						

REPRODUCE LOCALLY, include form number and edition date on a	all reproductions.	ORM APPROVED - OMB No. 0581-0055
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to det certificate is to be issued (7 U.S.C. 2 confidential until the certificate is issued.)	421). The information is held
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
F. 20 (027) 071770 01	OR EXPERIMENTAL NUMBER	
ENZA ZADEN BEHEER B.V.	14.2063	TELLMARK
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	6. FAX (include area code)
POSTBUS 7, 1600 AA ENKHUIZEN HALING 1º, 1602 DB ENKHUIZEN	+.31.228.315.844	+.31,228.315.854
HALING 1º 1602 DB ENKHVIZEN		7.3/,224,3.3,33/
THE NETHERLANDS	7. PVPO NUMBER 2006002	2 2 8
8. Does the applicant own all rights to the variety? Mark an "X" in the	ne appropriate block. If no, please expla	nin. YES NO
9. Is the applicant (individual or company) a U.S. national or a U.S. b	based company? If no, give name of c	ountry. YES NO
THE NETHERLANDS		النبقا السا
10. Is the applicant the original owner? YES	NO If no, please answer one	of the following:
a. If the original rights to variety were owned by individual(s), is YES b. If the original rights to variety were owned by a company(ies) YES	NO If no, give name of count	sed company?
11. Additional explanation on ownership (Trace ownership from origin	inal breeder to current owner. Use the re	everse for extra space if needed):
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not licens	sees) who meet the following criteria:	
 If the rights to the variety are owned by the original breeder, that penaltional of a country which affords similar protection to nationals of 	erson must be a U.S. national, national of the U.S. for the same genus and speci-	of a UPOV member country, or es.
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a c genus and species.	yed the original breeder(s), the company country which affords similar protection t	must be U.S. based, owned by o nationals of the U.S. for the same
3. If the applicant is an owner who is not the original owner, both the o	original owner and the applicant must m	eet one of the above criteria.
The original breeder/owner may be the individual or company who dir Act for definitions.	rected the final breeding. See Section 4	1(a)(2) of the Plant Variety Protection
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, a control number. The valid OMB control number for this information collection is 0581-0055, including the time for reviewing the instructions, searching existing data sources, gathering a	The time required to complete this information collect	tion is estimated to average 0.1 hour per response.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.

Form Approved OMB NO 9581-0055

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 5 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5964 (voice and TDD). USDA is an equal opportunity provider and employer.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MD 20705

EXHIBIT F
DECLARATION REGARDING DEPOSIT

NAME OF OWNER (S) ENZA ZADEN BEHEER B.V.	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country) POSTBUS 7, 1600 AA ENKHVIZEN HALING 1e, 1602 DB ENKHVIZEN	TEMPORARY OR EXPERIMENTAL DESIGNATION 14.2063
	THE NETHERLANDS	VARIETY NAME TELLMARK
NAME OF OWNER REPRESENTATIVE (S) MEL HOLLAND	ADDRESS (Street and No. or RD No., City, State, and Zip Code and Country) PO BOX 866, 525 LUCY BROWN LANE SAN JUAN BAUTISTA, CA 95045	PVPO TUMBER 6 0 0 2 2 8

I do hereby declare that during the life of the certificate a viable sample of propagating material of the subject variety will be deposited, and replenished as needed periodically, in a public repository in the United States in accordance with the regulations established by the Plant Variety Protection Office.

Signature

06/05/2006

Mond